

Long-range spin-qubit interaction mediated by microcavity polaritons

G. F. Quinteiro Rosen¹, J. Fernández-Rossier², and C. Piermarocchi¹

(1) *Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824 USA*

(2) *Department of Applied Physics, University of Alicante, Alicante, 03690 Spain*

(Dated: February 6, 2008)

We study the optically-induced coupling between spins mediated by polaritons in a planar micro-cavity. In the strong coupling regime, the vacuum Rabi splitting introduces anisotropies in the spin coupling. Moreover, due to their photon-like mass, polaritons provide an extremely long spin coupling range. This suggests the realization of two-qubit all-optical quantum operations within tens of picoseconds with spins localized as far as hundreds of nanometers apart.

Planar micro-cavities are semiconductor devices that confine the electromagnetic field by means of two parallel semiconductor mirrors. When a quantum well is placed inside a micro-cavity the optical excitations (excitons) in the well couple to the electromagnetic modes of the dielectric structure. In the so-called strong-coupling regime, excitons and cavity photons give rise to new states, cavity polaritons [1, 2], which appear in two branches separated by a vacuum Rabi splitting. In the long wavelength limit, and for a cavity exactly at resonance with the exciton energy, polaritons can be seen as hybrid states that are exactly half-matter and half-light. However, even with a half-matter character, polaritons have photon-like dispersive properties, determined by the reduced mass of the cavity-photon/exciton system. A photon confined in a $\lambda/2$ planar cavity has an effective mass of $M_\gamma = \hbar n \pi / c L$ (n is the refraction index and L is the length of the cavity) which is typically four to five orders of magnitude smaller than the exciton mass. The small polariton mass is known to affect the dynamics of optical excitations with phonons [3, 4], with interface disorder [5, 6], and it suggests the possibility of polariton Bose-Einstein condensates at room temperature. [7] In this paper, we show that the small polariton mass has also a strong effect on the mutual interaction between spins localized in a quantum well.

Proposals for quantum computers based on spin degrees of freedom require that individual qubits are placed close enough so to have a significant exchange interaction between them. This exchange interaction can be direct (i.e. induced by a controlled overlap of the wavefunctions), or indirect when mediated by spin excitations in a 2D electron gas [8] or by optical excitation across the semiconductor bandgap [9, 10, 11]. In the indirect schemes, the range of the spin coupling is related to the mass of the mediating particles, and the coupling decreases exponentially as a function of the distance between the spins. Here, we show that the small polariton mass gives an extremely long range for the spin coupling and introduces a non-exponential behavior. This implies that spin-qubits can be located several hundreds of nanometers apart while still retaining control on pair interaction through the use of polaritons.

We study two localized spins in a semiconductor quan-

tum well embedded in a planar micro-cavity. Our numerical results consider the case of two shallow neutral donors in GaAs (e. g. Si), but the theory is valid with minor modifications to other impurities and host semiconductors, as well as to charged quantum dots. Under strong coupling conditions, optically active excitons and intra-cavity photons combine into polaritons, while dark excitons remain unaffected. We consider a cavity excited by an extra-cavity continuous wave (cw) laser with frequency below the polariton resonance. In response to the laser, the cavity becomes polarized, without energy absorption, and this dynamic polarization is used to control the localized spins. Microscopically, the polarization is described in terms of a density of virtual polaritons, proportional to the laser intensity, that mediates an interaction between pairs of spins.

Our first step is to use the Hopfield canonical transformation [12] which allows us to treat the coupling between excitons and cavity-photons non-perturbatively. Due to spin-orbit and quantum confinement we can restrict our discussion to heavy-hole excitons only. Heavy-holes have a total spin component in the growth direction $j_z = \pm 3/2$. Optical excitations in the system are then described by lower polaritons with $J_z = \pm 1$, upper polaritons with $J_z = \pm 1$, and dark excitons with $J_z = \pm 2$. In the case of donors considered here the exciton-impurity spin interaction is given by an Heisenberg-like exchange interaction involving the electron in the exciton and the electron in the donor [11]. Therefore, if we restrict the excitation to σ_+ circularly polarized light, optically excited polaritons with $J_z = +1$ can only couple to states with $J_z = +2$ by scattering with the donors. Moreover, since we are considering microcavities in the strong coupling regime excited near the lower polariton resonance, we will neglect the upper polariton branch. This approximation will be further discussed below. The final Hamiltonian consists of a term H_0 describing lower polaritons with $J_z = +1$ and dark excitons with $J_z = +2$, a H_L term representing the coupling of the circularly polarized cw laser with the lower polariton at $k = 0$, and an interaction H_I describing polaritons and dark excitons

interacting with two localized impurities

$$\begin{aligned} H_0 &= \sum_k \Omega_k^P p_k^\dagger p_k + \Omega_k^X b_k^\dagger b_k \\ H_L &= \sqrt{\mathcal{A}} \phi_{1s} \mathcal{V} r_0 e^{i\omega_0 t} p_0 + h.c. \\ H_I &= \frac{J}{\mathcal{A}} \sum_{ikjk'\lambda} e^{i(k-k')R_\lambda} v_k v_{k'} H_{ij}^\lambda, \end{aligned}$$

where $\{i, j\}$ refer to the type of particle, either polariton (P) or dark exciton (X) with energy Ω_k^i and creation operators $p_k^\dagger$ and $b_k^\dagger$, respectively. The exciton energy is $\Omega_k^X = \epsilon_0 + \hbar^2 k^2 / 2M_X$ while the lower polariton energy is

$$\Omega_k^P = \frac{\Omega_k^X + \Omega_k^C}{2} - \sqrt{\frac{(\Omega_k^X - \Omega_k^C)^2}{4} + g_k^2}, \quad (1)$$

where $\Omega_k^C = \hbar c / n \sqrt{k^2 + (\pi/L)^2}$ is the energy of cavity photons, and g_k is the exciton-cavity photon coupling. We will consider microcavities at resonance, i.e. satisfying the condition $\Omega_0^C = \Omega_0^X$, with a vacuum Rabi splitting of $2g_0$. The index λ identifies the two localized impurities that we will call A and B . $\mathcal{V}$ and ω_0 are the Rabi energy and frequency of the exciting laser, and ϕ_{1s} is the excitonic enhancement in the light-matter coupling. Only the excitonic component of the polariton at $k = 0$ couples to the external laser field. This is taken into account by the Hopfield coefficient -which is the coefficient of proportionality between exciton and lower branch polariton operators- $r_0 = 1/\sqrt{2}$ [12]. At larger k the Hopfield coefficient approaches one, since in that limit the lower polariton becomes exciton-like. In order to take into account the finite range of the exchange interaction we use an exciton-impurity exchange interaction of the separable form $J v_k v_{k'}$ with $v_k = [1 + (\Lambda k)^2]^{-1}$ [13]. R_λ is the position of the impurity λ , and $\mathcal{A}$ is the quantization area. The spin dependent operators in H_I are defined as $H_{XX}^\lambda = \mathcal{S}_{XX}^\lambda b_{k'}^\dagger b_k$, $H_{XP}^\lambda = r_k \mathcal{S}_{XP}^\lambda b_{k'}^\dagger p_k$, $H_{PX}^\lambda = r_{k'}^* \mathcal{S}_{PX}^\lambda p_k^\dagger b_{k'}$, and $H_{PP}^\lambda = r_k^* r_{k'} \mathcal{S}_{PP}^\lambda p_k^\dagger p_{k'}$, where $\mathcal{S}_{XX}^\lambda = \mathcal{S}_{PP}^\lambda = s_z^\lambda s_z$, $\mathcal{S}_{PX}^\lambda = \mathcal{S}_{XP}^\lambda = s_x^\lambda s_x + s_y^\lambda s_y$, s^λ and s are electron spin operators in the shallow donor λ and in the polariton/exciton, respectively. The time dependent phase in H_L is eliminated by transforming the Hamiltonian in the rotating frame at frequency ω_0 .

We derive an effective Hamiltonian for the two electronic spins in the cavity due to the presence of the external laser field. This effective Hamiltonian can be written in terms of a level shift operator $R(\omega_0)$ [14] to second order in the laser coupling H_L as

$$H_{eff} = \mathcal{P} R(\omega_0) \mathcal{P} = \mathcal{P} H_L \frac{\mathcal{Q}}{\omega_0 - \mathcal{Q} H_0 \mathcal{Q} - \mathcal{Q} H_I \mathcal{Q}} H_L \mathcal{P} \quad (2)$$

where $\mathcal{Q}$ is the projector operator onto the subspace of one excitation (polariton or dark exciton) plus the two spins, and $\mathcal{P}$ is the projector operator onto the subspace of the two spins without excitations.

The next step is to make an expansion of the level shift operator operator in H_I keeping terms up to second order. Our result is therefore valid to second order both in the coupling with the external laser, $\mathcal{V}$, and in the exchange coupling J . However, since we are considering a microcavity in the strong coupling regime, we keep all orders in the exciton-cavity coupling by using the polariton picture. The theory can be expanded to include higher orders in $\mathcal{V}$ [15] and J [11]. In the case of spin 1/2, those contributions will modify the strength of the effects but will not change the general form and the qualitative behavior of the effective spin coupling.

The first order in the expansion of Eq. (2) in H_I is an effective magnetic field. It describes the Inverse Faraday effect of off-resonant circularly polarized radiation with frequency close to an optical resonance [16]. The Inverse Faraday effect on impurities in semiconductors was investigated in Refs. [11, 17], and was observed in Mn spins in quantum wells in Ref. [18]. This term disappears by using linearly polarized light because of time reversal symmetry. The second order in H_I plays the role of an effective interaction between the spins. We will focus our attention to this effective spin coupling, which is the one that can be used, for instance, to control spin entanglement and make optically controlled quantum gates.

The relevant terms in the level shift operator that contribute to the spin coupling are proportional to

$$H_{eff}^{(2)} \propto H_{PX}^A G_X^0 e^{i\phi} H_{XP}^B + H_{PP}^A G_P^0 e^{i\phi} H_{PP}^B + (A \rightleftharpoons B),$$

where $G_{P(X)}^0$ is the bare Green's operator for the polariton (exciton), the superscript (2) indicates the second order contribution in J only, and ϕ is the phase arising from the separation between impurities.

After some algebra, and using units of $\hbar = 1$, we obtain the final expression for the effective coupling as

$$H_{eff}^{(2)} = \frac{C}{\delta^2} [F_{RP} s_z^A s_z^B + F_{RX} (s_x^A s_x^B + s_y^A s_y^B)] \quad (3)$$

where $C = J^2 |\mathcal{V}|^2 |\phi_{1s}|^2 |r_0|^2$, and $\delta = \Omega_0^P - \omega_0$ is the laser-polariton detuning. The two functions F_{RP} and F_{RX} describe the polariton-mediated and the exciton-mediated contributions to the spin coupling and are defined as

$$\begin{aligned} F_{RP} &= \int_0^\infty \frac{dk}{2\pi} \frac{r_k^2 v_k^2 k J_0(kR)}{(\omega_0 - \Omega_k^P)} \\ F_{RX} &= \int_0^\infty \frac{dk}{2\pi} \frac{v_k^2 k J_0(kR)}{(\omega_0 - \Omega_k^X)}, \end{aligned}$$

where J_0 is the Bessel function of order zero and R is the inter-qubit separation. F_{RX} can be explicitly written in terms of modified Bessel functions K_0 and K_1 . In the large R limit the function F_{RX} has the 2D Yukawa form

$$F_{RX} \sim \frac{e^{-\sqrt{2M_X \delta} R}}{\sqrt{R}}. \quad (4)$$

Notice that the exponential behavior is characterized by a range $\ell \sim 1/\sqrt{M_X}$. The integral F_{R_P} has to be calculated by numerical quadrature.

Eq. (3) contains the main features of the polariton mediated spin coupling. The vacuum Rabi splitting resolves dark and optical active excitations, making the indirect interaction mediated by polaritons and dark excitons differ in strength and form. The overall interaction is spin-anisotropic. We can define a cut-off wave vector k_c such that for $k > k_c$ both $r_k \sim 1$ and $\Omega_k^P \sim \Omega_k^X$. The integrands of F_{R_P} and F_{R_X} coincide in that region since the polaritonic branch is exciton-like.

We can then rewrite

$$F_{R_P} \simeq F_{R_X} + \int_0^{k_c} (I_P - I_X) dk = F_{R_X} + \mathcal{D}_{PX}$$

where I_i is the integrand of either F_{R_P} or F_{R_X} . The term $\mathcal{D}_{PX}$ represents then the pure polariton contribution, while all the excitonic effects (dark-excitons plus polaritons at large k) are included in F_{R_X} . The cut-off k_c depends on the exciton cavity detuning and on the strength of the exciton-cavity coupling. Eq. (3) is rewritten as,

$$H_{eff} = \frac{C}{\delta^2} (F_{R_X} s^A \cdot s^B + \mathcal{D}_{PX} s_z^A s_z^B) .$$

Fig. 1 shows the relative strength of the Ising-like polariton-mediated and the isotropic exciton-mediated contributions for different values of the exciton-cavity coupling g_0 as a function of the spin distance. Excitonic atomic units are used, where energy is given in excitonic Ry^* and lengths are in Bohr radius a_B^* ; for GaAs, they correspond to $1Ry^* = 4.4meV$ and $a_B^* \sim 125\text{\AA}$. The analytical and numerical values for F_{R_X} , F_{R_P} and $\mathcal{D}_{PX}$ are presented in Fig. 2. Notice the existence of two distinct regions separated by a crossover distance R_c . For $R < R_c$, the dominant interaction has an isotropic Heisenberg form, while it changes to Ising-like for $R > R_c$. The first regime corresponds to an exciton-mediated coupling, while the second is a pure polariton-induced effect. Polaritons, due to their light mass, mediate a long range spin coupling interaction. However, since they have a fixed $J_z = +1$ they cannot flip the impurity spins, and can only induce an interaction diagonal in the impurity spin space. The upper polariton has also $J_z = +1$ and can only contribute to the Ising-like term. However, its contribution is reduced by $(\delta + 2g_0)^2$ due to the vacuum Rabi splitting $2g_0$. Fig. 3 shows that the strength of the exciton and polariton induced coupling in logarithmic scale. For the exciton, the interaction decays exponentially, while the polariton mediated term can survive up to extremely long spin-separations and shows a non-exponential behaviour. The long range nature of the polariton-mediated interaction presents important technological advantages for quantum information implementations. Using our parameters (see




FIG. 1: Relative weight of the pure polariton to exciton contribution as a function of the separation between the impurities. $g_0 = 0.4Ry^*$ (triangle); $g_0 = 0.25Ry^*$ (square); $g_0 = 0.1Ry^*$ (circle). In the plot we have used $\delta = 0.15Ry^*$, $\mathcal{V} = 0.16Ry^*$, $\Lambda = 0.25a_B^*$ and $J = 21Ry^*(a_B^*)^2$.




FIG. 2: Ising $J_{eff}^I = C/\delta^2 \mathcal{D}_{PX}$ (triangle) and Heisenberg $J_{eff}^X = C/\delta^2 F_{R_X}$ (square) contributions to the spin coupling. For completeness, the plot shows also the $J_{eff}^P = C/\delta^2 F_{R_P}$ (circle). The two regimes are separated by a crossover length $R_c \simeq 12 a_B^*$. $g_0 = 0.4Ry^*$, other parameters as in Fig. 1.

Fig. 1), we predict that the strength of the interaction is $J_{eff} \simeq 110^{-2}Ry^*$ for impurities separated by distances of the order of $R_c = 12 a_B^* \simeq 150nm$. An estimate for the time needed for an operation can be given as $T = \pi/J_{eff} \simeq 40ps$, which is much smaller than the typical dephasing time for impurity spin qubits. Recent measurements have reported a spin relaxation time of the order of μs for donors in GaAs [19]. To our knowledge, the spin decoherence time (T_2) of a single donor in GaAs has not been measured, but is also expected to be in the μs range. Moreover, notice that the time needed for a quantum operation does not change considerably when

we further increase the qubit separation. Even with an inter-qubit separation of 1 micron the time needed for one operation increases only by one order of magnitude to about 400 ps, and is still reasonably smaller than the decoherence time. With such a long range interaction, the realization of electric gates to control one-qubit operations and the use of localized magnetic field becomes feasible. The Ising-like interaction at long separation is not a limitation for quantum gate implementations. The polariton mediated coupling could be also used to control the nuclear spin of the donor in a scheme similar to the one in Ref. [8]. In contrast to other cavity QED-based quantum computing implementations [20], the scheme discussed here does not require 0D confined electromagnetic modes, which is much harder to achieve experimentally. In a planar cavity the lateral dimension is not limited by the optical wavelength, which provides a fully scalable geometry for the qubit.

A spin coupling can also be obtained by a real polariton population in a scheme analogue to the RKKY [21] spin coupling mechanism. The spin interaction induced by a 0D cavity and excitons in quantum dots has been recently investigated [22], and also in this 0D case the presence of a strong coupling generates anisotropies in the spin interaction. This approach is not appealing for quantum computing implementation since a real population of photocarriers will add decoherence to the spin-qubit. However, it would be interesting to explore the dynamics of spin in the presence of a dense polaritons population that condense in a phase coherent state, as observed recently in II-VI microcavities [23]. High-quality microcavities embedding Mn-doped magnetic quantum wells in the strong coupling regime have recently been realized [24]. The polariton mediated spin coupling could be explored in these systems as a method for the ultrafast control of the quantum well magnetization.

In conclusion, we have shown that the optical excitation of microcavity polaritons can couple spins localized in a quantum well. Due to the small polariton mass, the spin coupling has an extremely long range, and at large distances is Ising-like. The interaction is strong enough for the realization of quantum operations with spins located as far as several hundred of nanometers apart and within a time scale much shorter than the spin decoherence time. This interplay of polaritons and localized spins represents a peculiar feature of solid-state cavity QED, which has no equivalent in the atomic case.

This work was supported by NSF DMR-0312491 and by the Donald D. Harrington Foundation at the University of Texas Austin.




FIG. 3: Logarithmic plot of J_{eff}^X (triangle) and J_{eff}^P (circle), that shows the exponential decay and non-exponential behavior respectively ($g_0 = 0.4Ry^*$, other parameters as in Fig. 1.)

- [2] V. Savona, C. Piermarocchi, A. Quattropani, P. Schwendimann, and F. Tassone, *Phase Transitions* **68** 169 (1999).
- [3] G. Cassabois, A. L. C. Triques, F. Bogani, C. Delalande, Ph. Roussignol, and C. Piermarocchi, *Phys. Rev. B* **61**, 1696 (2000).
- [4] V. Savona and C. Piermarocchi, *phys. stat. sol. (a)* **164**, 45 (1997).
- [5] D. M. Whittaker, P. Kinsler, T. A. Fisher, M. S. Skolnick, A. Armitage, A. M. Afshar, M. D. Sturge, and J. S. Roberts, *Phys. Rev. Lett.* **77**, 4726 (1996).
- [6] V. Savona, C. Piermarocchi, A. Quattropani, F. Tassone, and P. Schwendimann, *Phys. Rev. Lett.* **78**, 4470 (1997).
- [7] H. Deng, G. Weihs, C. Santori, J. Bloch, and Y. Yamamoto, *Science* **298**, 199 (2002).
- [8] D. Mozyrsky, V. Privman, and M. L. Glasser, *Phys. Rev. Lett.* **86**, 5112 (2001).
- [9] C. Piermarocchi, Pochung Chen, L. J. Sham, and D. G. Steel, *Phys. Rev. Lett.* **86**, 167402 (2002).
- [10] G. Ramon, Y. Lyanda-Geller, T. L. Reinecke and L. J. Sham, *Phys. Rev. B* **71** 121305 (2005).
- [11] C. Piermarocchi and G. F. Quinteiro, *Phys. Rev. B* **70**, 235210 (2004).
- [12] J. J. Hopfield, *Phys. Rev.* **112**, 1555 (1958).
- [13] Y. Yamaguchi, *Phys. Rev.* **95** 1628 (1954).
- [14] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Atom-Photons Interactions*, (Chap. III) (Wiley New York, 1998).
- [15] J. Fernández-Rossier, C. Piermarocchi, P. Chen, A. H. MacDonald, and L. J. Sham, *Phys. Rev. Lett.* **93**, 127201 (2004).
- [16] L. D. Landau and E. M. Lifshitz, *Electrodynamics of continuous media*, (Chap. XI) (Pergamon Press, Oxford, Berlin 1984).
- [17] M. Combescot and O. Betbeder-Matibet, *Solid State Commun.* **132**, 129 (2004).
- [18] A. Gupta, R. Knobel, N. Samarth, and D. D. Awschalom, *Science* **292**, 2458 (2001).
- [19] C. Fu, C. Santori, C. Stanley, M. C. Holland, and Yoshi-

[1] C. Weisbuch, M. Nishioka, A. Ishikawa, and Y. Arakawa, *Phys. Rev. Lett.* **69**, 3314 (1992).

- hisa Yamamoto, Phys. Rev. Lett. **95**, 187405 (2005)
- [20] A. Imamoglu, D. D. Awschalom, G. Burkard, D. P. DiVincenzo, D. Loss, M. Sherwin, and A. Small, Phys. Rev. Lett. **83** 4204 (1999).
- [21] M. A. Ruderman and C. Kittel, Phys. Rev. **96**, 99 (1954); T. Kasuya, Progr. Theor. Phys. **16**, 45 (1956); K. Yosida, Phys. Rev. **106**, 893 (1957).
- [22] G. Chiappe, J. Fernández-Rossier, E. Louis, and E. V. Anda, Phys. Rev. B **72**, 245311 (2005).
- [23] M. Richard, J. Kasprzak, R. Romestain, R. André, and Le Si Dang, Phys. Rev. Lett. **94**, 187401 (2005).
- [24] S. Bahbah, R. Frey, R. André, G. Roosen, C. Flytzanis, Solid State Commun. **163** 147 (2005).